

MONOCLONAL ANTIBODY

For research use only, Not for diagnostic use.

Catalog No. NPM-M01

Anti Crumbs3 (Clone: 1B3-2D10)

BACKGROUND

Crumbs3 (Crb3) encodes the single transmembrane protein and is essential for normal development of epithelial tissues. Crb3 localizes at the apical plasma membrane and affects tight junction formation in polarized epithelial cells in vitro. Positive expression of Crb3 is detected in epithelial cells of the majority of hollow organs whereas stromal cells are negative in vertebrates. Enhanced expression of Crb3 promotes tumor cell migration and metastasis in a part of carcinomas. Crb3a and Crb3b, two isoforms of Crb3, are generated by alternative splicing. 1B3-2D10 monoclonal antibody was raised against the synthetic peptide miming the Crb3a-specific amino acid sequence.

Product type Primary antibody

Immunogen VGARVPPTPNLKLPPEERLI (Human Crb3a, Accession # NP_631900, Val101-ILe120)

 Rased in
 Mouse

 Clone number
 1B3-2D10

 Isotype
 IgG2a,k

Source Culture supernatant

Purification Cation exchange / gel filtration

Buffer Phosphate buffered saline containing 0.05 % ProClin300 as a preservative.

Concentration Indicated on the product label.

Volume 50 ug
Label Unlabeled

Specificity Human Crb3 isoform a

Reactivity human, mouse, porcine. Other species are not tested.

Store at 4 °C (avoid storing at -20 °C or lower).

Expiration date Indicated on the product label.

Application notes

Recommended dilutions

Immunohistochemistry (1~10 ug/mL).

Other applications are not covered under warranty.

Optimal dilutions/concentrations should be determined by the end user.

References

- Yokoyama Y., et al., (2022) Crumbs3 is expressed in oral squamous cell carcinomas and promotes cell migration and proliferation by affecting RhoA activity. Oncol Lett. 23(6):173. PMID: 35497937.
- lioka H., et al., (2019) Crumbs3 is a critical factor that regulates invasion and metastasis of colon adenocarcinoma via the specific interaction with FGFR1. Int J Cancer, 145(10):2740-2753. PMID: 30980524.

ANTIBODY CHARACTERIZATION




Fig.1 Immunohistochemical analysis of Crb3 in colon adenocarcinoma and adjacent non-neoplastic crypts in a patient tissue.




Fig.2 Immunohistochemical analysis of Crb3 in normal pig kidney.




Fig.3 Immunohistochemical analysis of Crb3 in the intestine of immunodeficient mice.

The optimal protocol for immunohistochemistry using anti-Crb3 monoclonal (clone # 1B3-2D10) should be determined by the end-user based on the following protocol.

- Specimen: Formalin fixed paraffin embedded (FFPE) tissue section.
- Antigen retrieval: Heat-Induced Epitope Retrieval (HIER) in 10 mM citrate buffer (pH 6.0)
- Primary antibody: anti-Crb3 monoclonal (clone # 1B3-2D10), 1~10 ug/ml. Incubate for 12 hr at 4 °C in a humidified chamber.
- Secondary antibody: Histofine simple stain MAX-PO(M) (Nichirei Bioscience #424132). Incubate for 0.5 hr at room temperature in a humidified chamber.
- Staining: TaKaRa DAB Substrate (TaKaRa Bio #MK210)

For research use only, Not for diagnostic use.



COSMO BIO CO., LTD.

[JAPAN]
TOYO EKIMAE BLDG. 2-20, TOYO 2-CHOME,
KOTO-KU. TOKYO 135-0016, JAPAN
Phone: +81-3-5632-9610
FAX: +81-3-5632-9619
URL: https://www.cosmobio.co.jp/



COSMO BIO USA

[Outside Japan] 2792 Loker Ave West, Suite 101 Carlsbad, CA 92010, USA email: info@cosmobiousa.com Phone/FAX: (+1) 760-431-4600 URL: www.cosmobiousa.com