




HSPB1 Antibody

| | |
|----------------------------|--|
| Product Code | CSB-RA010833A0HU |
| Abbreviation | Heat shock protein beta-1 |
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | P04792 |
| Immunogen | A synthesized peptide derived from human HSPB1 |
| Species Reactivity | Human |
| Tested Applications | ELISA, WB, IHC, IF, FC, IP; Recommended dilution: WB:1:500-1:5000, IHC:1:50-1:200, IF:1:20-1:200, IP:1:200-1:1000 |
| Relevance | Small heat shock protein which functions as a molecular chaperone probably maintaining denatured proteins in a folding-competent state (PubMed:10383393, PubMed:20178975). Plays a role in stress resistance and actin organization (PubMed:19166925). Through its molecular chaperone activity may regulate numerous biological processes including the phosphorylation and the axonal transport of neurofilament proteins (PubMed:23728742). |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Purification Method | Affinity-chromatography |
| Isotype | Rabbit IgG |
| Clonality | Monoclonal |
| Alias | Heat shock protein beta-1, HspB1, 28 kDa heat shock protein, Estrogen-regulated 24 kDa protein, Heat shock 27 kDa protein, HSP 27, Stress-responsive protein 27, SRP27, HSPB1, HSP27, HSP28 |
| Immunogen Species | Homo sapiens (Human) |
| Research Area | Signal Transduction |
| Gene Names | HSPB1 |
| Accession NO. | 10C11 |
| Image | |



Western Blot

Positive WB detected in: HeLa whole cell lysate, PC3 whole cell lysate, 293T whole cell lysate, LO2 whole cell lysate, HT-29 whole cell lysate, K562 whole cell lysate


All lanes: Hsp27 antibody at 0.62µg/ml

Secondary


Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 23 KDa


Observed band size: 27 KDa




IHC image of CSB-RA010833A0HU diluted at 1:61.9 and staining in paraffin-embedded human pancreatic tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC image of CSB-RA010833A0HU diluted at 1:61.9 and staining in paraffin-embedded human cervical cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of MCF-7 cells with CSB-RA010833A0HU at 1:20, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L).




Immunoprecipitating Hsp27 in HeLa whole cell lysate

Lane 1: Rabbit control IgG instead of CSB-RA010833A0HU in HeLa whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000)

Lane 2: CSB-RA010833A0HU (3µg) + HeLa whole cell lysate (500µg)

Lane 3: HeLa whole cell lysate (20µg)



Overlay histogram showing HeLa cells stained with CSB-RA010833A0HU (red line) at 1:50. The cells were fixed with 70% Ethylalcohol (18h) and then permeabilized with 0.3% Triton X-100 for 2 min. The cells were then incubated in 1x PBS /10% normal goat serum to block non-specific protein-protein interactions followed by primary antibody for 1 h at 4°C. The secondary antibody used was FITC goat anti-rabbit IgG (H+L) at 1/200 dilution for 1 h at 4°C. Control antibody (green line) was used under the same conditions. Acquisition of >10,000 events was performed.

Description

The product is an HSPB1 recombinant monoclonal antibody. Its production process includes the cloning of the human HSPB1 DNA gene into the vector and subsequent transfection of the clones into the cell line for in vitro expression. This HSPB1 antibody is purified using affinity-chromatography. It shows reactivity with the human HSPB1 protein. And it has been validated in multiple applications, including ELISA, WB, IHC, IF, FC, and IP.

HSPB1, an anti-apoptotic protein, is over-expressed in various tumors and promotes adverse outcomes. Overexpression of HSPB1 is frequently related to increased resistance to radiotherapy and to anti-cancer drugs, such as cisplatin, doxorubicin, and etoposide. HSPB1 has emerged as a major therapeutic target in cancer therapy due to its cytoprotective and oncogenic properties.