

🕜 Tel: +1-301-363-4651 🛛 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com 🍙

MAP3K3 Antibody

Product CodeCSB-RA013425A0HUAbbreviationMitogen-activated protein kinase kinase kinase 3StorageUpon receipt, store at -20°C or -80°C. Avoid repeated freeze.Uniprot No.Q99759ImmunogenA synthesized peptide derived from human MAP3K3Species ReactivityHumanTested ApplicationsELISARelevanceComponent of a protein kinase signal transduction cascade. Mediates activation of the NF-kappa-B, AP1 and DDIT3 transcriptional regulators.FormLiquidConjugateNon-conjugatedStorage BufferRabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Purification MethodAffinity-chromatographyIsotypeRabbit IgGClonalityMonoclonalAliasMitogen-activated protein kinase kinase kinase 3, MAPK/ERK kinase kinase 3, MEKK inase 3, MEKK 3, MAP3K3, MAPKKK3, MEKK3Research AreaSignal TransductionGene NamesMAP3K3Accession NO.2H11		
StorageUpon receipt, store at -20°C or -80°C. Avoid repeated freeze.Uniprot No.Q99759ImmunogenA synthesized peptide derived from human MAP3K3Species ReactivityHumanTested ApplicationsELISARelevanceComponent of a protein kinase signal transduction cascade. Mediates activation of the NF-kappa-B, AP1 and DDIT3 transcriptional regulators.FormLiquidConjugateNon-conjugatedStorage BufferRabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Purification MethodAffinity-chromatographyIsotypeRabbit IgGClonalityMonoclonalAliasMitogen-activated protein kinase kinase kinase 3, MAPK/ERK kinase kinase 3, MEK kinase 3, MEKK 3, MAP3K3, MAPKKK3, MEKK3Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionGene NamesMAP3K3	Product Code	CSB-RA013425A0HU
Uniprot No.Q99759ImmunogenA synthesized peptide derived from human MAP3K3Species ReactivityHumanTested ApplicationsELISARelevanceComponent of a protein kinase signal transduction cascade. Mediates activation of the NF-kappa-B, AP1 and DDIT3 transcriptional regulators.FormLiquidConjugateNon-conjugatedStorage BufferRabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Purification MethodAffinity-chromatographyIsotypeRabbit IgGClonalityMonoclonalAliasMitogen-activated protein kinase kinase si, MAPK/ERK kinase kinase 3, MEK kinase 3, MEKK 3, MAP3K3, MAPKKK3, MEKK3Immunogen SpeciesHomo sapiens (Human)Research AreaSignal Transduction MAP3K3	Abbreviation	Mitogen-activated protein kinase kinase kinase 3
ImmunogenA synthesized peptide derived from human MAP3K3Species ReactivityHumanTested ApplicationsELISARelevanceComponent of a protein kinase signal transduction cascade. Mediates activation of the NF-kappa-B, AP1 and DDIT3 transcriptional regulators.FormLiquidConjugateNon-conjugatedStorage BufferRabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Purification MethodAffinity-chromatographyIsotypeRabbit IgGClonalityMonoclonalAliasMitogen-activated protein kinase kinase kinase 3, MAPK/ERK kinase kinase 3, MEK kinase 3, MEKK 3, MAP3K3, MAPKKK3, MEKK3Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionMereMAP3K3	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Species ReactivityHumanTested ApplicationsELISARelevanceComponent of a protein kinase signal transduction cascade. Mediates activation of the NF-kappa-B, AP1 and DDIT3 transcriptional regulators.FormLiquidConjugateNon-conjugatedStorage BufferRabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Purification MethodAffinity-chromatographyIsotypeRabbit IgGClonalityMonoclonalAliasMitogen-activated protein kinase kinase kinase 3, MAPK/ERK kinase kinase 3, MEK kinase 3, MEKK 3, MAP3K3, MAPKKK3, MEKK3Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionMAP3K3MAP3K3	Uniprot No.	Q99759
Tested ApplicationsELISARelevanceComponent of a protein kinase signal transduction cascade. Mediates activation of the NF-kappa-B, AP1 and DDIT3 transcriptional regulators.FormLiquidConjugateNon-conjugatedStorage BufferRabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Purification MethodAffinity-chromatographyIsotypeRabbit IgGClonalityMonoclonalAliasMitogen-activated protein kinase kinase si, MAPK/ERK kinase kinase 3, MEK kinase 3, MEKK 3, MAP3K3, MAPKKK3, MEKK3Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionGene NamesMAP3K3	Immunogen	A synthesized peptide derived from human MAP3K3
RelevanceComponent of a protein kinase signal transduction cascade. Mediates activation of the NF-kappa-B, AP1 and DDIT3 transcriptional regulators.FormLiquidConjugateNon-conjugatedStorage BufferRabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Purification MethodAffinity-chromatographyIsotypeRabbit IgGClonalityMonoclonalAliasMitogen-activated protein kinase kinase kinase 3, MAPK/ERK kinase 3, MEK kinase 3, MEKK 3, MAP3K3, MAPKKK3, MEKK3Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionGene NamesMAP3K3	Species Reactivity	Human
of the NF-kappa-B, AP1 and DDIT3 transcriptional regulators.FormLiquidConjugateNon-conjugatedStorage BufferRabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Purification MethodAffinity-chromatographyIsotypeRabbit IgGClonalityMonoclonalAliasMitogen-activated protein kinase kinase kinase 3, MAPK/ERK kinase kinase 3, MEK kinase 3, MEKK 3, MAP3K3, MAPKKK3, MEKK3Research AreaSignal TransductionGene NamesMAP3K3	Tested Applications	ELISA
ConjugateNon-conjugatedStorage BufferRabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Purification MethodAffinity-chromatographyIsotypeRabbit IgGClonalityMonoclonalAliasMitogen-activated protein kinase kinase kinase 3, MAPK/ERK kinase kinase 3, MEK kinase 3, MEKK 3, MAP3K3, MAPKKK3, MEKK3Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionGene NamesMAP3K3	Relevance	
Storage BufferRabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Purification MethodAffinity-chromatographyIsotypeRabbit IgGClonalityMonoclonalAliasMitogen-activated protein kinase kinase 3, MAPK/ERK kinase kinase 3, MEK kinase 3, MEKK 3, MAP3K3, MAPKKK3, MEKK3Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionGene NamesMAP3K3	Form	Liquid
azide and 50% glycerol.Purification MethodAffinity-chromatographyIsotypeRabbit IgGClonalityMonoclonalAliasMitogen-activated protein kinase kinase 3, MAPK/ERK kinase kinase 3, MEK kinase 3, MEKK 3, MAP3K3, MAPKKK3, MEKK3Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionGene NamesMAP3K3	Conjugate	Non-conjugated
IsotypeRabbit IgGClonalityMonoclonalAliasMitogen-activated protein kinase kinase kinase 3, MAPK/ERK kinase kinase 3, MEK kinase 3, MEKK 3, MAP3K3, MAPKKK3, MEKK3Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionGene NamesMAP3K3	Storage Buffer	
ClonalityMonoclonalAliasMitogen-activated protein kinase kinase kinase 3, MAPK/ERK kinase kinase 3, MEK kinase 3, MEKK 3, MAP3K3, MAPKKK3, MEKK3Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionGene NamesMAP3K3	Purification Method	Affinity-chromatography
AliasMitogen-activated protein kinase kinase kinase 3, MAPK/ERK kinase kinase 3, MEK kinase 3, MEKK 3, MAP3K3, MAPKKK3, MEKK3Immunogen SpeciesHomo sapiens (Human)Research AreaSignal TransductionGene NamesMAP3K3	Isotype	Rabbit IgG
MEK kinase 3, MEKK 3, MAP3K3, MAPKKK3, MEKK3 Immunogen Species Homo sapiens (Human) Research Area Signal Transduction Gene Names MAP3K3	Clonality	Monoclonal
Research AreaSignal TransductionGene NamesMAP3K3	Alias	5 1 <i>7 7</i>
Gene Names MAP3K3	Immunogen Species	Homo sapiens (Human)
	Research Area	Signal Transduction
Accession NO. 2H11	Gene Names	MAP3K3
	Accession NO.	2H11

1