




# GABRA5 Antibody

<b>Product Code</b>	CSB-RA559038A0HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	P31644
<b>Immunogen</b>	A synthesized peptide derived from human GABA A Receptor alpha 5
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA, WB, IHC; Recommended dilution: WB:1:500-1:5000, IHC:1:50-1:200
<b>Relevance</b>	GABA, the major inhibitory neurotransmitter in the vertebrate brain, mediates neuronal inhibition by binding to the GABA/benzodiazepine receptor and opening an integral chloride channel.
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Purification Method</b>	Affinity-chromatography
<b>Isotype</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Product Type</b>	Recombinant Antibody
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Neuroscience
<b>Gene Names</b>	GABRA5
<b>Accession NO.</b>	5C11

## Image



### Western Blot

Positive WB detected in: Ntera-2 whole cell lysate, HEK293 whole cell lysate


All lanes: GABA A Receptor alpha 5 antibody at 1:1000

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 53 kDa

Observed band size: 70 kDa



IHC image of CSB-RA559038A0HU diluted at 1:100 and staining in paraffin-embedded human brain tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.

## Description

The GABRA5 monoclonal antibody's DNA sequence was inserted into the plasmid, which was subsequently transfected into the cell line for expression. The GABRA5 recombinant monoclonal antibody was produced after purification using affinity chromatography. This rabbit IgG GABRA5 recombinant antibody has been evaluated in scientific applications such as ELISA, WB, and IHC. It exclusively responds with GABRA5 from humans.

GABRA5 is abundantly expressed in the hippocampus, mainly in the extrasynaptic area of CA1 pyramidal cells, where it regulates tonic inhibitory conductance and may cause synaptic plasticity and memory impairments. GABRA5-mediated phasic inhibition via vasoactive intestinal peptide (VIP) input to interneurons is important for anxiety modulation while the GABRA5 tonic suppression through this subunit may regulate spatial learning.