




# TRAF2 Antibody

<b>Product Code</b>	CSB-RA786226A0HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	Q12933
<b>Immunogen</b>	A synthesized peptide derived from human TRAF2
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA, WB, IHC, FC, IP; Recommended dilution: WB:1:500-1:5000, IHC:1:50-1:200, FC:1:20-1:200, IP:1:200-1:1000
<b>Relevance</b>	Regulates activation of NF-kappa-B and JNK and plays a central role in the regulation of cell survival and apoptosis. Required for normal antibody isotype switching from IgM to IgG. Has E3 ubiquitin-protein ligase activity and promotes 'Lys-63'-linked ubiquitination of target proteins, such as BIRC3, RIPK1 and TICAM1. Is an essential constituent of several E3 ubiquitin-protein ligase complexes, where it promotes the ubiquitination of target proteins by bringing them into contact with other E3 ubiquitin ligases. Regulates BIRC2 and BIRC3 protein levels by inhibiting their autoubiquitination and subsequent degradation; this does not depend on the TRAF2 RING-type zinc finger domain. Plays a role in mediating activation of NF-kappa-B by EIF2AK2/PKR. In complex with BIRC2 or BIRC3, promotes ubiquitination of IKBKE.
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Purification Method</b>	Affinity-chromatography
<b>Isotype</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Product Type</b>	Recombinant Antibody
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Cancer; Cardiovascular; Cell biology; Signal transduction
<b>Gene Names</b>	TRAF2
<b>Accession NO.</b>	9A5

## Image



**Western Blot**

Positive WB detected in: HepG2 whole cell lysate, HeLa whole cell lysate, 293T whole cell lysate, Jurkat whole cell lysate, Raji whole cell lysate, K562 whole cell lysate


All lanes: TRAF2 antibody at 1:1500

Secondary


Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 56, 62, 55, 54 kDa


Observed band size: 56 kDa




IHC image of CSB-RA786226A0HU diluted at 1:100 and staining in paraffin-embedded human kidney tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



IHC image of CSB-RA786226A0HU diluted at 1:100 and staining in paraffin-embedded human cervical cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit IgG polymer labeled by HRP and visualized using 0.05% DAB.



Overlay histogram showing HeLa cells stained with CSB-RA786226A0HU (red line) at 1:50. The cells were fixed with 70% Ethylalcohol (18h) and then incubated in 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1µg/1\*10<sup>6</sup>cells) for 1 h at 4°C. The secondary antibody used was FITC-conjugated goat anti-rabbit IgG (H+L) at 1/200 dilution for 30min at 4°C. Control antibody (green line) was Rabbit IgG (1µg/1\*10<sup>6</sup>cells) used under the same conditions. Acquisition of >10,000 events was performed.



Immunoprecipitating TRAF2 in HeLa whole cell lysate

Lane 1: Rabbit control IgG instead of CSB-RA786226A0HU in HeLa whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000)

Lane 2: CSB-RA786226A0HU(2μg)+ HeLa whole cell lysate(500μg)

Lane 3: HeLa whole cell lysate (10μg)

## Description

TRAF2 is an important adaptor protein that transduces signals after ligating to certain receptors including those binding TNF and is also an E3 ubiquitin ligase. TRAF2 is required for TNF- $\alpha$ -mediated activation of MAPK8/JNK and NF- $\kappa$ B. Studies of TRAF2-deficient mice showed that TRAF2 plays an essential role in mediating cell survival, normal adaptive immune responses, and lymphocyte homeostasis. TRAF2 promotes survival, proliferation, and metastasis of tumor cells through the NF- $\kappa$ B pathway by directly interacting with various TNF receptors.

Compared with the polyclonal and monoclonal antibodies of TRAF2, this TRAF2 recombinant antibody has the features of increased reproducibility and control, animal-free technology, high degree of monovalency, high batch-to-batch consistency, easier isotype conversion, etc. And it has been validated in ELISA, WB, IHC, FC, IP.