




GBA Antibody

| | |
|----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Product Code | CSB-RA869334A0HU |
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | P04062 |
| Immunogen | A synthesized peptide derived from human GBA |
| Species Reactivity | Human |
| Tested Applications | ELISA, WB; Recommended dilution: WB:1:500-1:5000 |
| Relevance | extracellular exosome, lysosomal lumen, lysosomal membrane, glucosylceramidase activity, receptor binding, cellular response to tumor necrosis factor, ceramide biosynthetic process, glucosylceramide catabolic process, glycosphingolipid metabolic process, negative regulation of inflammatory response |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Purification Method | Affinity-chromatography |
| Isotype | Rabbit IgG |
| Clonality | Monoclonal |
| Product Type | Recombinant Antibody |
| Immunogen Species | Homo sapiens (Human) |
| Research Area | Neuroscience; Cancer; Metabolism; Signal transduction |
| Gene Names | GBA |
| Accession NO. | 4H4 |

Image



Western Blot

Positive WB detected in: HeLa whole cell lysate, MCF-7 whole cell lysate, 293T whole cell lysate, U87 whole cell lysate

All lanes: GBA antibody at 1:2000

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 60, 58, 55, 51, 30 kDa

Observed band size: 60 kDa

Description

GBA encodes the glucocerebrosidase (GCCase), a lysosomal glycoside



hydrolase that catalyzes the hydrolysis of the glycolipid glucosylceramide (GlcCer) to glucose and ceramide. It is involved in the metabolism of sphingolipids. GBA deficiency results in the buildup of GlcCer and disruption of lipid balance. Gaucher's disease has been linked to human GBA gene mutations, which are numerically the most important risk factor for developing Parkinson's disease (PD), accounting for at least 5% of all PD cases. Furthermore, sporadic PD brains show a decrease in GBA activity.

The recombinant GBA antibody was generated in vitro through inserting cloned GBA genes into expression vectors. The expression vector was then inserted into a mammalian cell to express this GBA antibody. It has been validated in ELISA, WB. Every step in the production was controlled strictly. You have no worries about the quality.