

POLYCLONAL ANTIBODY

# Anti-SLC9A7 (NHE7) (Human) pAb

Code No.  
BMP043

Quantity  
50 µL

Form  
Affinity Purified

**BACKGROUND:** SLC9A7, also known as (Na<sup>+</sup>, K<sup>+</sup>)/H<sup>+</sup> exchanger 7 (NHE7), locates predominantly in the trans-Golgi network (TGN) and plays a role in organellar pH homeostasis. By using the yeast two-hybrid system, the secretory carrier membrane protein 2 (SCAMP2) is revealed to be associated with SLC9A7 as a binding partner, and this indicates the involvement of SCAMP2 in the recruitment of SLC9A7 to TGN.

**SOURCE:** This antibody was affinity purified from rabbit serum. The rabbit was immunized with a synthetic peptide derived from human SLC9A7.

**FORMULATION:** 50 µL volume of PBS containing 50% glycerol, pH 7.2. No preservative is contained.

**STORAGE:** This antibody solution is stable for one year from the date of purchase when stored at -20°C.

**REACTIVITY:** This antibody can be used to stain endogenous antigen in paraffin embedded human tissues including the liver and cerebellum by Immunohistochemistry. The reactivity has been confirmed by Western blotting, Immunocytochemistry and intracellular Flow cytometry to detect the full length of human SLC9A7 transiently expressed in HEK 293T cells.

## APPLICATIONS:

Western blotting; 1:1,000 for chemiluminescence detection system

Immunoprecipitation; Not tested

Immunohistochemistry; 1:1,000

Heat treatment is necessary for staining paraffin embedded sections.

Autoclave; 125°C for 5 minutes in 10 mM citrate buffer containing 0.05% Tween-20 (pH 6.0).

Immunocytochemistry; 1:1,000

Flow cytometry; 1:1,000 (final concentration)

Detailed procedure is provided in the following **PROTOCOLS**.

## INTENDED USE:

For Research Use Only. Not for use in diagnostic procedures.


## SPECIES CROSS REACTIVITY:

Species	Human	Mouse	Rat
Tissues	liver, cerebellum	Not tested	Not tested
Reactivity on IHC	+		


## REFERENCES:

- 1) Milosavljevic, N., *et al.*, *Cell Rep.* **7**, 689-696 (2014) [IC, IHC]
- 2) Kagami, T., *et al.*, *Mol. Membr. Biol.* **25**, 436-447 (2008)
- 3) Lin, P. J. C., *et al.*, *J. Cell Sci.* **118**, 1885-1897 (2005)
- 4) Numata, M., and Orłowski, J., *J. Biol. Chem.* **276**, 17387-17394 (2001)

**cerebellum**



**liver**



**Immunohistochemical detection of SLC9A7 on paraffin embedded section of human cerebellum and liver with BMP043. Multi pathological types tissue array (MBL) was used for this application.**

## PROTOCOLS:

Immunohistochemical staining for paraffin-embedded sections

- 1) Deparaffinize the sections with Xylene 3 times for 3-5 minutes each.


- 2) Wash the slides with Ethanol 3 times for 3-5 minutes each.
- 3) Wash the slides with PBS 3 times for 3-5 minutes each.
- 4) Heat treatment

Heat treatment by Autoclave:

Heat the slides immersed in retrieval solution [10 mM citrate buffer containing 0.05% Tween-20 (pH 6.0)] at 125°C for 5 minutes in pressure boiler. After boiling, the slides should remain in the pressure boiler until the temperature is cooled down to 80°C. Let the immersed slides further cool down at room temperature for 40 minutes.

- 5) Remove the slides from the retrieval solution and cover each section with 0.3% H<sub>2</sub>O<sub>2</sub> in PBS for 15 minutes at room temperature to block endogenous peroxidase activity. Wash 3 times in PBS containing 0.05% Tween-20 for 5 minutes each.
  - 6) Remove the slides from PBS, wipe gently around each section and cover tissues with blocking buffer (0.5% BSA and 5% Normal goat serum in PBS) for 30 minutes at room temperature to block non-specific staining. Do not wash.
  - 7) Tip off the blocking buffer, wipe gently around each section and cover tissues with primary antibody diluted with blocking buffer as suggested in the **APPLICATIONS**.
- Note:** It is essential for every laboratory to determine the optional titers of the primary antibody to obtain the best result.
- 8) Incubate the sections for 2 hours at room temperature.
  - 9) Wash the slides 3 times in PBS for 5 minutes each.
  - 10) Wipe gently around each section and cover tissues with ENVISION/HRP polymer reagent (DAKO; code no. K1491). Incubate for 15 minutes at room temperature. Wash as in step 9).
  - 11) Visualize by reacting for 5 minutes with DAB substrate solution (DAKO; code no. K3465). \*DAB is a suspect carcinogen and must be handled with care. Always wear gloves.
  - 12) Wash the slides in water for 5 minutes.
  - 13) Counter stain in hematoxylin for 1 minute, wash the slides 3 times in water for 5 minutes each, and then immerse the slides in PBS for 5 minutes. Dehydrate by immersing in Ethanol 3 times for 3 minutes each, followed by immersing in Xylene 3 times for 3 minutes each.
  - 14) Now ready for mounting.


(Positive control for Immunohistochemistry; liver, cerebellum)



**Immunocytochemical detection of SLC9A7 in 293T transiently expressing SLC9A7 with BMP043 (right) or normal rabbit IgG (left).**

### Immunocytochemistry

- 1) Culture the cells at an appropriate condition on a glass slide. (for example, spread 1x10<sup>4</sup> cells for one slide, then incubate in a CO<sub>2</sub> incubator for one night.)
- 2) Wash the cells 3 times with PBS.
- 3) Fix the cells by immersing the slide in PBS containing 4% paraformaldehyde (PFA) for 15 minutes at 4°C.
- 4) Wash the slide 2 times with PBS containing 0.5% BSA.
- 5) Immerse the slide in PBS containing 0.1% Triton X-100 for 15 minutes at room temperature.
- 6) Wash the cells 2 times with PBS containing 0.5% BSA, 0.1% Triton X-100.
- 7) Immerse the slide in blocking buffer (0.1% Triton X-100, 0.5% BSA, 5% Normal goat serum, 0.1 mg/mL of human IgG in PBS) for 60 minutes at room temperature.
- 8) Wash the slide 2 times with PBS containing 0.5% BSA, 0.1% Triton X-100.
- 9) Tip off the washing buffer, add the primary antibody diluted with blocking buffer at a titer as suggested in the **APPLICATIONS** onto the cells and incubate for 1 hour at room temperature (Optimizations of antibody titer or incubation condition is recommended if necessary.)
- 10) Wash the slide 3 times with PBS containing 0.5% FCS, 0.1% Triton X-100.
- 11) Add 100 µL of PE conjugated anti-rabbit IgG (Beckman Coulter; code no. 732743) at a titer of 1:200 diluted with blocking buffer. Incubate in the dark at room temperature for 30 minutes.
- 12) Wash the slide 3 times with PBS containing 0.5% BSA, 0.1% Triton X-100.
- 13) Wipe excess liquid from slide but take care not to touch the cells. Never leave the cells to dry.
- 14) Promptly add mounting medium onto the slide, then put a cover slip on it.




**Flow cytometric analysis of intracellular SLC9A7 expression on 293T transiently expressing SLC9A7 and Azami green\*. The staining intensity of BMP043 is shown in the vertical axis with Azami Green fluorescence on the horizontal axis.**

### Flow cytometric analysis

We usually use Fisher tubes or equivalents as reaction tubes for all steps described below.

- 1) Wash the cells 3 times with PBS containing 2% FCS.

- 2) Resuspend the cells with PBS containing 2% FCS ( $5 \times 10^6$  cells/mL).
  - 3) Add 50  $\mu$ L of the cell suspension into each tube, and centrifuge at 500 x g for 1 minute at room temperature. Remove supernatant by careful aspiration.
  - 4) Add 100  $\mu$ L of 4% paraformaldehyde (PFA) in PBS to the cell pellet after tapping. Mix well, then fix the cells for 15 minutes at 4°C.
  - 5) Wash the cells 2 times with PBS containing 0.5% BSA.
  - 6) Add 100  $\mu$ L of PBS containing 0.1% Triton X-100 to the cell pellet after tapping. Mix well, then permeabilize the cells for 15 minutes at room temperature (20~25°C).
  - 7) Wash the cells 2 times with PBS containing 0.5% BSA, 0.1% Triton X-100..
  - 8) Add 20  $\mu$ L of blocking buffer (PBS containing 0.1% Triton X-100, 0.5% BSA, 5% normal goat serum, 0.1 mg/mL human IgG) to the cell pellet after tapping. Mix well and incubate for 60 minutes at 4°C.
  - 9) Add 20  $\mu$ L of the primary antibody at a titer as suggested in the **APPLICATIONS** diluted with blocking buffer. Mix well and incubate for 1 hour at room temperature.
  - 10) Wash the cells 3 times with PBS containing 0.5% BSA, 0.1% triton X-100.
  - 11) Add 20  $\mu$ L of PE conjugated anti-rabbit IgG at a titer of 1:200 (Beckman Coulter; code no. 732743) diluted with blocking buffer. Mix well and incubate in the dark for 30 minutes at room temperature.
  - 12) Wash the cells 3 times with PBS containing 0.5% BSA, 0.1% triton X-100.
  - 13) Resuspend the cells with 500  $\mu$ L of PBS containing 2% FCS, analyze by a flow cytometer.
- 3) Mix the sample with equal volume of Laemmli's sample buffer.
  - 4) Incubate the samples for 1 hour at 37°C and centrifuge at 10,000 x g for 5 minutes. Transfer the supernatant into a new tube. Load 10  $\mu$ L of sample per lane on a 1-mm-thick SDS-polyacrylamide gel and carry out electrophoresis.
  - 5) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at 1 mA/cm<sup>2</sup> for 1 hour in a semi-dry transfer system (Transfer Buffer: 25 mM Tris, 190 mM glycine, 20% MeOH). See the manufacturer's manual for precise transfer procedure.
  - 6) To reduce nonspecific binding, soak the membrane in 5% skimmed milk (in PBS, pH 7.2) for 2 hours at room temperature, or overnight at 4°C.
  - 7) Incubate the membrane for 2 hours at room temperature with primary antibody diluted with PBS (pH 7.2) containing 2% skimmed milk as suggested in the **APPLICATIONS**. (The concentration of antibody will depend on the conditions.)
  - 8) Wash the membrane with PBS-T [0.05% Tween-20 in PBS] (5 minutes x 3 times).
  - 9) Incubate the membrane with 1:2,000 of Anti-IgG (Rabbit) pAb-HRP (MBL; code no. 458) diluted with 2% skimmed milk (in PBS, pH 7.2) for 1 hour at room temperature.
  - 10) Wash the membrane with PBS-T (10 minutes x 3 times).
  - 11) Drain excess buffer on the membrane, and incubate membrane with an appropriate chemiluminescence reagent for 1 minute.
  - 12) Remove extra reagent from the membrane by dabbing with a paper towel, and seal it in plastic wrap.
  - 13) Expose and develop the film under usual settings. The condition for exposure and development may vary.



**Western blot analysis of SLC9A7 expression in Myc-tagged SLC9A7 transfected 293T (2, 3) and parental cell (1) using BMP043 (1, 2) or anti-Myc-tag antibody (3, MBL; code no. M047-3).**

#### **RELATED PRODUCTS:**

- BMP023 Anti-SLC9A1 (NHE1) (Human) pAb (polyclonal)
- BMP075 Anti-SLC9A2 (NHE2) (Human) pAb (polyclonal)
- BMP049 Anti-SLC9A6 (NHE6) (Human) pAb (polyclonal)
- BMP043 Anti-SLC9A7 (NHE7) (Human) pAb (polyclonal)
- BMP061 Anti-SLC9A9 (NHE9) (Human) pAb (polyclonal)
- PM035 Normal Rabbit IgG

#### **SDS-PAGE & Western Blotting**

- 1) Wash cells (approximately  $2 \times 10^6$  cells) 3 times with PBS and suspend with 100  $\mu$ L of cold Lysis buffer (10 mM Tris-HCl pH 7.5, 150 mM NaCl, 1% Triton X-100, 1% Sodium deoxycholate, 0.1% SDS) containing appropriate protease inhibitors. Incubate it at 4°C with rotating for 30 minutes, then sonicate briefly (up to 10 seconds).
- 2) Centrifuge the tube at 12,000 x g for 10 minutes at 4°C and transfer the supernatant to another fresh tube.