

For Research Use Only.
Not for use in diagnostic procedures.



Anti-CLOCK (Mouse) mAb

CODE No. D334-3

CLONALITY Monoclonal
CLONE CLNT1
ISOTYPE Mouse IgG1 κ
QUANTITY 100 μ L, 1 mg/mL

SOURCE Purified IgG from hybridoma supernatant
IMMUNOGEN N-terminal region (Met¹-Gly¹²⁰) of mouse Clock
FORMURATION PBS containing 50% Glycerol (pH 7.2). No preservative is contained.
STORAGE This antibody solution is stable for one year from the date of purchase when stored at -20°C.

APPLICATION-CONFIRMED

Western blotting 1-5 μ g/mL for chemiluminescence detection system

APPLICATION-REPORTED

Immunoprecipitation Reference 1) and 2)

SPECIES CROSS REACTIVITY on WB

Species	Human	Mouse	Rat	Hamster
Sample	Not tested	Liver nuclear extract	Not tested	Not tested
Reactivity		+		

Entrez Gene ID 12753 (Mouse)

REFERENCES
1) Yoshitane, H., *et al.*, *EMBO Rep.* **13**, 455-461 (2012) [IP]
2) Yoshitane, H., *et al.*, *Mol Cell Biol.* **29**, 3675-3686 (2009) [WB, IP]

For more information, please visit our web site <http://ruo.mbl.co.jp/>



RELATED PRODUCTS

Antibodies

D334-3	Anti-CLOCK (Mouse) mAb (CLNT1)
D333-3	Anti-CLOCK (Mouse) mAb (CLSP3)
D335-3	Anti-BMAL1 (Mouse) mAb (B1BH2)
CY-P1016	Anti-SIRT1 pAb
RN032P	Anti-CIRBP pAb

Kits

CY-1151	CycLex [®] SIRT1/Sir2 Deacetylase Fluorometric Assay Kit
CY-1152	CycLex [®] SIRT2 Deacetylase Fluorometric Assay Kit
CY-1153	CycLex [®] SIRT3 Deacetylase Fluorometric Assay Kit
CY-1156	CycLex [®] SIRT6 Deacetylase Fluorometric Assay Kit


Recombinant proteins (Human, Active)

CY-E1151	NAD ⁺ -Dependent Deacetylase SIRT1
CY-E1152	NAD ⁺ -Dependent Deacetylase SIRT2
CY-E1153	NAD ⁺ -Dependent Deacetylase SIRT3
CY-E1156	NAD ⁺ -Dependent Deacetylase SIRT6

SDS-PAGE & Western blotting

- 1) Mix 10 μ L of Mouse liver nuclear extract with 10 μ L of Laemmli's sample buffer.
- 2) Boil the sample for 3 min. and centrifuge. Load 20 μ L of the sample per lane in a 1-mm-thick SDS-polyacrylamide gel (7.5% acrylamide) for electrophoresis.
- 3) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at 300 mA for 1 hr. in a wet transfer system (Transfer Buffer: 25 mM Tris, 190 mM glycine, 20% MeOH). See the manufacturer's manual for precise transfer procedure.
- 4) To reduce nonspecific binding, soak the membrane in 1% skimmed milk (in TBS, pH 7.2) for 1 hr. at room temperature.
- 5) Incubate the membrane with primary antibody diluted with 1% skimmed milk (in TBS, pH 7.2) as suggested in the **APPLICATION** for 1 hr. at 37°C, 2 hr. at room temperature or overnight at 4°C. (The concentration of antibody will depend on the conditions.)
- 6) Wash the membrane 3 times for 2 min., 5 min. and 10 min. each with 1% skimmed milk (in TBS, pH 7.2).
- 7) Incubate the membrane with the 1:5,000 anti-IgG (Mouse) pAb-HRP (MBL; code no. 330) diluted with 1% skimmed milk (in TBS, pH 7.2) for 2 hr. at room temperature or overnight at 4°C.
- 8) Wash the membrane 3 times for 2 min., 5 min. and 10 min. each with TBS-T [0.05% Tween-20 in TBS].
- 9) Wash the membrane 1 time for 2 min. with TBS.
- 10) Wipe excess buffer on the membrane, and then incubate it with appropriate chemiluminescence reagent for 1 min. Remove extra reagent from the membrane by dabbing with paper towel, and seal it in plastic wrap.
- 11) Expose to an X-ray film in a dark room for 3 min. Develop the film as usual. The condition for exposure and development may vary.

(Positive controls for Western blotting; Mouse liver nuclear extracts)



*The heavy/light chains derived from IgG in the samples.
(These bands are detected depending on a sample.)

Western blot analysis of mouse CLOCK from liver nuclear extracts

- 1: ZT6 (zeitgeber time; 6 h)
- 2: ZT18 (zeitgeber time; 18 h)

Immunoblotted with Anti-CLOCK (Mouse) mAb (D334-3)

Data were provided by Mr. Kentaro Hirose, Dr. Hikari Yoshitane, Ph.D. and Dr. Yoshitaka Fukada, Ph.D. (Department of Biophysics and Biochemistry, Graduate School of Science, University of Tokyo)