

For Research Use Only.
Not for use in diagnostic procedures.

Smart-IP Series

Mouse IgG2a (isotype control) -Magnetic Beads

CODE No.	M076-11
CLONALITY	Monoclonal
CLONE	6H3
ISOTYPE	Mouse IgG2a κ
QUANTITY	20 tests (Slurry: 1 mL)
SOURCE	Purified IgG from hybridoma supernatant
IMMUNOGEN	KLH
REACTIVITY	No specific binding is detected on immunoprecipitation.
FORMULATION	Covalently antibody conjugated 10 mg magnetic beads in 1 mL PBS/0.1% BSA/0.09% NaN ₃ . *Azide may react with copper or lead in plumbing system to form explosive metal azides. Therefore, always flush plenty of water when disposing materials containing azide into drain.
STORAGE	This beads suspension is stable for one year from the date of purchase when stored at 4°C.
APPLICATION-CONFIRMED	
<u>Immunoprecipitation</u>	50 μ L of beads slurry/sample


RELATED PRODUCTS

For more information, please visit our web site <https://ruo.mbl.co.jp/>.

The descriptions of the following protocols are examples. Each user should determine the appropriate condition.

Immunoprecipitation

- 1) Wash 2×10^6 cells 3 times with PBS and suspend them in 500 μ L of cold Extraction buffer [50 mM Tris-HCl (pH 7.5), 150 mM NaCl, 1% NP-40].
- 2) Centrifuge the tube at 12,000 x g for 5 min. at 4°C and transfer the supernatant to another tube.
- 3) Add magnetic beads as suggested in the **APPLICATION** into 500 μ L of the cell lysate. Mix well and incubate with gentle agitation for 30 min. at 4°C.
- 4) Place the tube on the magnetic rack (MBL; code no. 3190) for a few seconds.
- 5) Remove the supernatant.
- 6) Add 1 mL of cold Wash buffer [50 mM Tris-HCl (pH 7.5), 150 mM NaCl, 0.05% NP-40] and resuspend the magnetic beads.
- 7) Place the tube on the magnetic rack for a few seconds.
- 8) Remove the supernatant.
- 9) Repeat Steps 6)-8) 3 times.
- 10) Resuspend the magnetic beads in 20 μ L of Laemmli's sample buffer, boil for 3 min., and place the tube on the magnetic rack for a few seconds.
- 11) Load 20 μ L of the sample per lane in a 1-mm-thick SDS-polyacrylamide gel and carry out electrophoresis.
- 12) Visualize the protein bands by CBB staining.



Immunoprecipitation of DDDDK-tagged protein

Sample: HEK293T cell lysate from 2×10^6 cells + DDDDK-tagged GFP 10 μ g

Lane 1: Input (10 μ L/lane)

Lane 2: Post-IP beads of Mouse IgG2a (isotype control)-Magnetic Beads (M076-11)

Lane 3: Post-IP beads of Anti-DDDDK-tag mAb-Magnetic Beads (M185-11)