

Anti-DDDDK-tag mAb-Alexa Fluor[®] 488

CODE No.	M185-A48
CLONALITY	Monoclonal
CLONE	FLA-1
ISOTYPE	Mouse IgG2a κ
QUANTITY	100 μ L, 1 mg/mL
SOURCE	Purified IgG from hybridoma supernatant
IMMUNOGEN	KLH conjugated DYKDDDDK peptide
REACTIVITY	This antibody reacts with N-terminal, Internal and C-terminal DDDDK-tagged (DYKDDDDK) proteins.
FORMULATION	PBS containing 1% BSA and 0.1% ProClin 150.
STORAGE	This antibody solution is stable for one year from the date of purchase when stored at 4°C.

APPLICATIONS-CONFIRMED

<u>Immunocytochemistry</u>	0.5-1 μ g/mL
<u>Flow cytometry</u>	0.5 μ g/mL

REFERENCES	1) Kasubuchi, M., <i>et al.</i> , <i>Sci. Rep.</i> 7 , 5168 (2017) [IC] 2) Kobayashi, Y., <i>et al.</i> , <i>PLoS Pathog.</i> 12 , e1005785 (2016) [IC]
-------------------	--

RELATED PRODUCTS

For more information, please visit our web site <https://ruo.mbl.co.jp/>.

LABEL LICENSES:


This product is provided under an agreement between LIFE TECHNOLOGIES Corporation, and Medical & Biological Laboratories Co., LTD.

Alexa Fluor[®] is a registered trademark of Molecular Probes, Inc.

The descriptions of the following protocols are examples. Each user should determine the appropriate condition.

Immunocytochemistry

- 1) Spread the cells in the nutrient condition on a glass slide, then incubate in a CO₂ incubator overnight.
- 2) Remove the culture supernatant by careful aspiration.
- 3) Fix the cells by immersing the slide in 4% paraformaldehyde (PFA)/PBS for 10 min. at room temperature (20~25°C).
- 4) Prepare a wash container such as a 500 mL beaker with a magnetic stirrer. Then wash the fixed cells on the glass slide by soaking the slide with a plenty of PBS in the wash container for 5 min. Take care not to touch the cells. Repeat another wash once more.
- 5) Immerse the slide in 0.2% Triton X-100/PBS for 10 min. at room temperature.
- 6) Wash the slide in a plenty of PBS as in the step 4).
- 7) Cover each cell with Clear Back (human Fc receptor blocking reagent, MBL, code no. MTG-001) for 5 min. at room temperature.
- 8) Add 200 µL of the primary antibody diluted with PBS as suggested in the **APPLICATIONS** onto the cells and incubate for 1 hr. at room temperature. (Optimization of antibody concentration or incubation condition is recommended if necessary.)
- 9) Wash the slide in a plenty of PBS as in the step 4).
- 10) Wipe excess liquid from slide but take care not to touch the cells. Never leave the cells to dry.
- 11) Counter stain with DAPI for 5 min. at room temperature.
- 12) Wash the slide in a plenty of PBS as in the step 4).
- 13) Promptly add mounting medium onto the slide, then put a cover slip on it.





Immunocytochemical detection of DDDDK-tagged protein in HeLa

Green: M185-A48
Blue: DAPI

Flow cytometric analysis for adherent cells

- 1) Detach the cells from culture dish.
- 2) Wash the cells once with 1 mL of washing buffer [PBS containing 2% fetal calf serum (FCS)].
- 3) Resuspend the cells with washing buffer (3×10^6 cells/mL).
- 4) Add 100 μ L of the cell suspension into each tube, and centrifuge at 500 x g for 1 min. at room temperature (20~25°C). Remove supernatant by careful aspiration.
- 5) Add 20 μ L of Clear Back (human Fc receptor blocking reagent, MBL, code no. MTG-001) to the cell pellet after tapping. Mix well and incubate for 5 min. at room temperature.
- 6) Add 40 μ L of the primary antibody at the concentration as suggested in the **APPLICATIONS** diluted in the washing buffer. Mix well and incubate for 30 min. at room temperature.
- 7) Add 1 mL of the washing buffer followed by centrifugation at 500 x g for 1 min. at room temperature. Remove supernatant by careful aspiration. Repeat another wash once more.
- 8) Resuspend the cells with 500 μ L of the washing buffer and analyze by a flow cytometer.



Flow cytometric detection of DDDDK-tagged protein in HeLa

Open: M185-A48

Closed: Isotype control (MBL, code no. M076-A48)

Upper: DDDDK-tagged protein in HeLa

Lower: Parental cell (HeLa)