For Research Use Only. Not for use in diagnostic procedures.



Anti-mouse IL-33

(Functional Grade)

CODE No. M188-3

CLONALITY Monoclonal

CLONE 2C7

 $\begin{array}{ll} \textbf{ISOTYPE} & \textbf{Mouse IgG2a} \; \kappa \\ \textbf{QUANTITY} & 200 \; \mu L, \; 1 \; \text{mg/mL} \end{array}$

SOURCE Purified IgG from hybridoma supernatant **IMMUNOGEN** Mouse IL-33, 109-266 aa (recombinant)

FORMURATION 1 mg/mL in PBS. Azide free, 0.22 μm sterile-filtered

Endotoxin level is <0.5 EU/mg antibody, as determined by the LAL assay

STORAGE This antibody solution is stable for one year from the date of purchase when stored at 4°C.

APPLICATIONS-CONFIRMED

Western blotting 1 μg/mL for chemiluminescence detection system

APPLICATIONS-REPORTED

Neutralization Reference 1)

SPECIES CROSS REACTIVITY on WB

| Species | Human | Mouse | Rat | Hamster |
|------------|--------------|--------------|------------|------------|
| Cells | Transfectant | Transfectant | Not tested | Not Tested |
| Reactivity | - | + | | |

Entrez Gene ID 77125 (Mouse)

REFERENCES 1) Ohno, T., et al., PLoS One 6, e18404 (2011) [Neutralization]

2) Matsuyama, Y., et al., J. Rheumatol. 37, 18-25 (2010)

3) Smith, D. E., et al., Clin. Exp. Allergy 40, 200-208 (2010)

4) Cayrol, C. and Girard, J. P., PNAS. 106, 9021-9026 (2009)

5) Luthi, A. U., et al., Immunity 31, 84-98 (2009)

6) Schmitz, J., et al., Immunity 23, 479-490 (2005)

For more information, please visit our web site https://ruo.mbl.co.jp/



RELATED PRODUCTS

Antibody M187-3 anti-Mouse IL-33 (1F11) [WB, Nu] M161-3 anti-Mouse IL-33 (4G4) [WB, IP] M138-3 anti-Human IL-33 (5H1) [WB, IP] anti-Human IL-33 (polyclonal) [WB, IHC] PM033 D065-3 anti-Human ST2 (HB12) [WB, IP, FCM, IHC] anti-Human ST2 (FB9) [WB, IP, FCM, ELISA] D066-3 anti-Human ST2 (2A5) [WB, IP, FCM] D067-3 D067-4 anti-Human ST2-FITC (2A5) D067-5 anti-Human ST2-PE (2A5)

Isotype control antibody

M075-3M2 Mouse IgG1 isotype control (2E12) [FCM, FA] M076-3M2 Mouse IgG2a isotype control(6H3) [FCM, FA]

ELISA Kit

7650 Human IL-33 Cytokine domain Detection Kit 5332 Ab-Match ASSEMBLY mouse IL-33 ELISA

7638 ST2 ELISA kit

Recombinant Protein

B005-10 Human IL-33

Others


8455 MaxBlot

WB: Western blotting
IP: Immunoprecipitation
IHC: Immunohistochemistry
FCM: Flow cytometry
Nu: Neutralization
FA: Functional Analysis

SDS-PAGE & Western blotting

- 1) Wash 1 x 10⁷ cells 3 times with PBS and suspends them in 1 mL of Laemmli's sample buffer, then sonicate briefly (up to 10 seconds).
- 2) Boil the samples for 3 minutes and centrifuge. Load 10 μL of the sample per lane in a 1-mm-thick SDS-polyacrylamide gel (12.5% acrylamide) for electrophoresis.
- 3) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at 1 mA/cm² for 1 hour in a semi-dry transfer system (Transfer Buffer: 25 mM Tris, 190 mM glycine, 20% MeOH). See the manufacturer's manual for precise transfer procedure.
- 4) To reduce nonspecific binding, soak the membrane in 10% skimmed milk (in PBS, pH 7.2) for overnight at 4°C.
- 5) Wash the membrane with PBS-T [0.05% Tween-20 in PBS] (5 minutes x 3 times).
- 6) Incubate the membrane with primary antibody diluted with MaxBlot Solution 1 (MBL; code no. 8455) as suggested in the **APPLICATIONS** for 1 hour at room temperature. (The concentration of antibody will depend on the conditions.)
- 7) Wash the membrane with PBS-T (5 minutes x 3 times).
- 8) Incubate the membrane with the 1:10,000 HRP-conjugated anti-mouse IgG (MBL; code no. 330) diluted with 1% skimmed milk (in PBS, pH 7.2) for 1 hour at room temperature.
- 9) Wash the membrane with PBS-T (5 minutes x 3 times).
- 10) Wipe excess buffer on the membrane, and then incubate it with appropriate chemiluminescence reagent for 1 minute. Remove extra reagent from the membrane by dabbing with paper towel, and seal it in plastic wrap.
- 11) Expose to an X-ray film in a dark room for 2 minutes. Develop the film as usual. The condition for exposure and development may vary.

(Positive control for Western blotting; mouse IL-33 trasnfectant)



Western blot analysis of mouse IL-33

Lane 1: mouse IL-33/293T Lane 2: 293T

Immunoblotted with M188-3 in MaxBlot solution 1