

POLYCLONAL ANTIBODY

ER Marker

Anti-Calnexin pAb

Code No.
PM060

Quantity
100 µL

Form
Affinity Purified

BACKGROUND: The endoplasmic reticulum (ER) is a eukaryotic organelle, which serves many general functions, including the facilitation of protein folding. Calnexin is a 90 kDa integral membrane protein of the ER. Calnexin is one of the chaperone proteins, which play a major role in the quality control of the ER by the retention of incorrectly folded proteins.

SOURCE: This antibody was purified from rabbit serum using affinity column. The rabbit was immunized with the synthetic peptide corresponding to N-terminus of human Calnexin.

FORMULATION: 100 µL volume of PBS containing 50% glycerol, pH 7.2. Contains no preservative.

STORAGE: This antibody solution is stable for one year from the date of purchase when stored at -20°C.

REACTIVITY: This antibody reacts with human Calnexin for Western blotting, Immunoprecipitation and Immunocytochemistry.

APPLICATIONS:

Western blotting; 1:1,000

Immunoprecipitation; 2 µL/300 µL of cell extract from 1 x 10⁷ cells

Immunohistochemistry; Not tested

Immunocytochemistry; 1:1,000

Flow cytometry; Not tested

Detailed procedures are provided in the following **PROTOCOLS**.

SPECIES CROSS REACTIVITY:

Species	Human	Mouse	Rat
Cells	HeLa, 293T, A549, Jurkat	NIH/3T3	Not Tested
Reactivity on WB	+	-	


REFERENCES:

- 1) Kleizen, B., and Braakman, I., *Curr. Opin. Cell Biol.* **16**, 343-349 (2004)
- 2) David, V., *et al.*, *J. Biol. Chem.* **268**, 9585-9592 (1993)

INTENDED USE:

For Research Use Only. Not for use in diagnostic procedures.

The descriptions of the following protocols are examples. Each user should determine the appropriate condition.



Western blotting analysis of Calnexin expression on HeLa (1), 293T (2), A549 (3) and Jurkat (4) using PM060.


PROTOCOLS:

SDS-PAGE & Western blotting

- 1) Wash cells (approximately 1 x 10⁷ cells) 3 times with PBS and resuspend them in 1 mL of Laemmli's sample buffer.
- 2) Boil the samples for 2 minutes and centrifuge. Load 10 µL of sample per lane on a 1-mm-thick SDS-polyacrylamide gel and carry out electrophoresis.
- 3) Blot the protein to a polyvinylidene difluoride (PVDF) membrane at 1 mA/cm² for 1 hour in a semi-dry transfer system (Transfer Buffer: 25 mM Tris, 190 mM glycine, 20% methanol). See the manufacturer's manual for precise transfer procedure.
- 4) To reduce nonspecific binding, soak the membrane in 10% skimmed milk (in PBS, pH 7.2) for 1 hour at room temperature, or overnight at 4°C.
- 5) Incubate the membrane for 1 hour at room temperature with primary antibody diluted with 1% skimmed milk (in PBS, pH 7.2) as suggested in the **APPLICATIONS**. (The concentration of antibody will depend on the conditions.)
- 6) Wash the membrane with PBS-T [0.05% Tween-20 in PBS] (5 minutes x 3).
- 7) Incubate the membrane with 1:10,000 HRP-conjugated anti-rabbit IgG (MBL, code no. 458) diluted with 1% skimmed milk (in PBS, pH 7.2) for 1 hour at room temperature.
- 8) Wash the membrane with PBS-T (5 minutes x 3).

- 9) Wipe excess buffer off the membrane, and incubate membrane with an appropriate chemiluminescence reagent for 1 minute.
- 10) Remove extra reagent from the membrane by dabbing with a paper towel, and seal it in plastic wrap.
- 11) Expose the membrane onto an X-ray film in a dark room for 3 minutes. Develop the film under usual settings. The conditions for exposure and development may vary.

(Positive controls for Western blotting; HeLa, 293T, A549, Jurkat)



Immunoprecipitation of Calnexin from HeLa with normal rabbit IgG (1) or PM060 (2). After immunoprecipitated with the antibody, immunocomplex was resolved on SDS-PAGE and immunoblotted with PM060.

Immunoprecipitation


- 1) Wash cells (approximately 1×10^7 cells) 3 times with PBS and resuspend them in 1 mL of cold Lysis buffer [50 mM Tris-HCl (pH 7.5), 150 mM NaCl, 0.05% NP-40] containing protease inhibitors at appropriate concentrations. Incubate it at 4°C with rotating for 30 minutes; thereafter, briefly sonicate the mixture (up to 10 seconds).
- 2) Centrifuge the tube at 12,000 x g for 10 minutes at 4°C and transfer the supernatant to another tube.
- 3) Add primary antibody as suggested in the **APPLICATIONS** into 300 μ L of the supernatant. Mix well and incubate with gentle agitation for 30-120 minutes at 4°C. Add 20 μ L of 50% protein A agarose beads resuspended in the cold Lysis buffer. Mix well and incubate with gentle agitation for 60 minutes at 4°C.
- 4) Centrifuge the tube at 2,500 x g for 10 seconds, and carefully discard the supernatant using a pipettor without disturbing the beads.
- 5) Resuspend the beads with cold Lysis buffer.
- 6) Centrifuge the tube at 2,500 x g for 10 seconds, and carefully discard the supernatant.
- 7) Repeat steps 5)-6) 3-5 times
- 8) Resuspend the beads in 20 μ L of Laemmli's sample buffer, boil for 3-5 minutes, and centrifuge for 5 minutes. Use 10 μ L/lane for the SDS-PAGE analysis.
(See **SDS-PAGE & Western blotting**.)

(Positive control for Immunoprecipitation; HeLa)

Immunocytochemistry

- 1) Culture the cells in the appropriate condition on a glass slide. (for example, spread 1×10^4 cells for one slide, then incubate in a CO₂ incubator for one night.)
- 2) Wash the glass slide 2 times with PBS.
- 3) Fix the cells by immersing the slide in PBS containing 4% paraformaldehyde for 10 minutes at room temperature.
- 4) Wash the glass slide 3 times with PBS.
- 5) Immerse the slide in PBS containing 0.2% Triton X-100 for 10 minutes at room temperature.
- 6) Wash the glass slide 2 times with PBS.
- 7) Add the primary antibody diluted with 2% FCS/PBS as suggested in the **APPLICATIONS** onto the cells and incubate for 1 hour at room temperature (Optimization of antibody concentration or incubation condition are recommended if necessary).
- 8) Wash the glass slide 2 times with PBS.
- 9) Add 100 μ L of 1:500 Alexa Fluor[®] 488 conjugated anti-rabbit IgG (Thermo Fisher Scientific, code no. A110374) diluted with PBS onto the cells. Incubate for 30 minutes at room temperature. Keep out light by aluminum foil.
- 10) Wash the glass slide 3 times with PBS.
- 11) Counter stain with DAPI for 5 minutes at room temperature.
- 12) Wash the glass slide 2 times with PBS.
- 13) Wipe excess liquid off the slide but take care not to touch the cells. Never leave the cells to dry.
- 14) Promptly add mounting medium onto the slide, then put a cover slip on it.

(Positive control for Immunocytochemistry; A549)



Immunocytochemical detection of Calnexin in A549 with PM060.
Green: Anti-Calnexin pAb (MBL, code no. PM060)
Blue: DAPI counter stain

RELATED PRODUCTS:

Please visit our website at <https://ruo.mbl.co.jp/>.